





# Infection Related GN

Tariq Zayan

Nephrologist Sur Hospital, Oman


# Introduction

#### Immune-Mediated

IRGN is glomerulonephritis due to an immunologic kidney response caused by a nonrenal infection.

# The disease's epidemiology, morphology and progression are significantly influenced by:

- The timing of the infection
- The patient's age
- The pathogen involved
- Comorbid conditions
- Socioeconomic conditions
- Healthcare access
- Changing patterns in infectious diseases



# Types of IRGN



Occur after the infection has resolved.

#### Parainfectious Forms

Occur during an active, often persistent infection lasting weeks. (increasing importance in older patients).

# **Epidemiology Misnomer**

# Streptococcal Infections

In modern times it accounts for only 28%- 47% of postinfectious acute GN.

#### Staphylococcus Infections

aureus or epidermidis: isolated in 12%-24% of cases.

# Gram-negative bacteria

Isolated in up to 22% of cases

#### **Global Incidence**

Is difficult to determine, as it is often transient and is likely underreported or unrecognized.

#### **Incidence and Prevalence**

#### **Decline in Developed Countries**

The incidence of acute post-streptococcal glomerulonephritis (APSGN), historically the most common form of IRGN, has seen a marked decline in developed countries.

#### **WHO Estimates**

WHO estimates approximately 470,000 new cases of APSGN annually worldwide, with 97% occurring in low-socioeconomic regions.

#### **Incidence Rates**

- In developing countries, the median incidence of APSGN is reported at 24.3 cases per 100,000 person-years among children.
- In affluent communities, this figure drops to 6.2 cases per 100,000 person-years.

# **Changing Demographics**

#### **Shifting Profile**

Prevalence of *Staph* RGN is higher than acute poststreptococcal GN in geographic areas with adequate resources.

# Increase in Adult Cases

Particularly among older adults with comorbidities such as diabetes and hypertension...etc

#### **Comorbidities**

Approximately **1/3** adults with postinfectious glomerulonephritis have at least one comorbid condition

### **Prognosis**



Children

Generally, have favorable outcomes with complete recovery of kidney function post-infection.



**Healthy Adults** 

Also tend to recover well but may experience more complications if underlying health issues are present.



#### **Older Adults & Comorbidities**

Often face a more guarded prognosis due to chronic disease.
Studies indicate that adults with comorbid factors may experience more chronic kidney injury following

**IRGN** 

# **Pathophysiology**

#### In Situ Formation

Positively charged circulating antigens cross the negatively charged basement membrane and bind to antigen-specific antibodies, forming subepithelial immune complexes.

# **Circulating Immune Complex Deposition**

Antigens already bound to antibodies in circulation that deposit in glomeruli.

#### **Plasmin Binding**

Antigens deposited in glomeruli can bind plasmin, activating immunological processes such as:

- Complement activation (via alternative or classical pathways)
- Recruitment of inflammatory cells, particularly neutrophils.

# Superantigens & Neutrophil extracellular traps (NETs)

#### Superantigens

produced by pathogens like Staph. & Chlamydia pneumoniae → polyclonal T-cell activation, → release of pro-inflammatory cytokines such as IL-2, IFN-γ, and TNF-α. → autoantibodies and immune complexes may deposit in the glomeruli, triggering complement activation and inflammatory damage.

# Excessive or dysregulated NET formation contributes to glomerular injury by:

- Inducing endothelial damage.
- Acting as a scaffold for immune complex deposition.
- Triggering complement activation (via the alternative or lectin pathway).
- Exposing autoantigens (e.g., histones or MPO), promoting autoimmune responses.

# Genetic Susceptibility and Predisposition

#### The Major Histocompatibility Complex (MHC) region

Specific HLA types, such as HLA-DRB1\\*03011, have been found at higher frequencies in patients with acute poststreptococcal glomerulonephritis (APSGN) compared to healthy individuals

# Pathogenesis of Post-infectious/Poststreptococcal glomerulonephritis (PIGN)

#### Key nephritogenic antigens include:

- Nephritis-associated plasmin receptor (GAPDH).
- Streptococcal pyrogenic exotoxin B (SPEB).
- They activate the alternate complement pathway, resulting in hypocomplementemia
- · Have an affinity towards plasmin and glomerular proteins.

# Clinical and Diagnostic Features

Clinically

Include acute nephritic syndrome, transient arterial hypertension (50%), or nephrotic syndrome in 5–10% of cases.

Serologically

C3 consumption occurs in ~90% of cases and resolves within 6-8 weeks.

Anti-streptococcal antibodies

(e.g., antistreptolysin O, DNase B) are key diagnostic markers.

- Possible biomarkers:
  - Nephritis-associated plasmin receptor, (streptococcal glyceraldehyde-3-phosphate dehydrogenase)
  - Streptococcal pyrogenic exotoxin B (SPE-B)

## Histopathological Characteristics:

1

2

3

#### Early stages

Diffuse global intra-/endocapillary proliferation with neutrophilic infiltration (exudative GN). Occasional crescents.

#### Immunofluorescence

typically reveals C3-dominant or codominant glomerular staining

#### Later stages

Mesangial expansion and hypercellularity may precede full resolution.

<sup>1.</sup> Nasr SH, et al KI 2013.

<sup>2.</sup> Kanjanabuch T et al Nature Reviews Nephrology.2009

#### **IRGN** in Adults

#### IRGN in adults

differs significantly, becoming more parainfectious due to persistent infections.

#### Management

prioritizes treating the underlying infection.

#### **Common Infections**

#### Sites:

Respiratory tract, skin, urinary tract, and systemic infections like sepsis.

#### **Etiology:**

Predominantly Staphylococcus species, gram-negative bacteria, and fewer Streptococcus cases.

# Histological Patterns

Acute phase

Diffuse intracapillary proliferation and neutrophil-rich exudates.

Chronic progression

Mesangioproliferative or membranoproliferative patterns.

Immunohistochemistry and Electron Microscopy

**Typical findings:** C3-dominant or co-dominant patterns.

Acute phase deposits: Subepithelial "humps."

Chronic phase: Predominantly mesangial deposits.

# IgA- dominant IRGN

#### **Pathogens**

MRSA, MSSA, E. coli, S. epidermidis, & Klebsiella bacteraemia in patients with underlying comorbidities, especially diabetes

#### Presentation

- Bacteremia is often, but not always, found, although the presentation may be delayed.
- Reported in patients with skin and joint infections, pneumonia, osteomyelitis, and endocarditis

#### Hypocomplementemia

May be seen in 30%-50% of cases.

#### **Key Features**

IgA-dominant immune response with strong C3 reactivity

#### **Differential Diagnosis**

IgA nephropathy and ANCA-associated vasculitis due to overlapping serological and histological findings.

#### **Kidney Biopsy**

- A kidney biopsy is useful in adults, particularly when the culture evidence of infection is elusive, or the diagnosis is in doubt.
- In some cases, a biopsy may be critical at arriving at the correct diagnosis, as comorbidities (DM, HIV) contribute to the difficulty in making the diagnosis

#### Persistent Low C3

In classical postinfectious GN, a persistently low C3 in serum beyond 12weeks may be an indication for kidney biopsy to exclude C3 glomerulonephritis (C3GN).

#### Infective Endocarditis-Associated IRGN

#### **Prosthetic Valves**

Prosthetic valves or structural heart valve lesions

#### **Risk Factors**

Substance abuse, Elderly, DM, HCV, HIV or Immunocompromised

#### **Pathogens**

Strep. viridans (low virulence, left heart) and Staph. aureus (high virulence, right heart).

#### **Compatible IE History**

- ECHO with vegetations,
   Fever, new or changed
   cardiac murmur
- Splenomegaly and other classical features

#### **Blood Cultures**

- + ve blood cultures(90-98%)
- Negative in 2-10%.

#### **Histological Findings**

Extracapillary proliferative GN with or without necrosis.

# **Shunt Nephritis**

# Associated with ventriculoatrial shunts

- Associated with ventriculoatrial shunts colonized by Streptococcus epidermidis.
- Lower risk with ventricular-jugular shunts.
- Least with ventricular-peritoneal shunts.
- May present after months or years following shunt placement & sometimes after shunt revision

#### **Symptoms and Diagnosis**

- Symptoms may be non-specific
- ANCA titers may be positive
- Presents as membranoproliferative GN with prominent IgM deposits.

#### **Treatment of Bacterial IR-GN**

# Treat the underlying infection

- No RCT to guide treatment of any of the 4 infection-related GNs
- Treat the underlying infection with antibiotics directed against the organism
- Although this will not alter the course- it can prevent the spread of infection within the community

#### **Manage Symptoms**

- Treat oedema,
   hypertension- as well as persistent proteinuria
- May require the support of KRT

#### Glucocorticoids and Immunosuppression

The utility of glucocorticoids and immunosuppression is unproven and carries serious potential risk - even in cases with crescentic GN

#### **HIV- related GN**

#### **Global Problem**

As it contributing significantly to CKD

#### Biopsy Importance

Should be performed, when feasible, to evaluate the morphology-multiple pathologies that occurs

#### **Outcome Factors**

- Persistence of viral replication
- Response to ART
- (APOL1) Genetic predisposition
- Coinfection with other viruses
- Development of IC disease or TMA

#### **ART Initiation**

Current consensus data from the

#### TEMPRANO study,

demonstrate benefit of ART initiation at diagnosis - regardless of CD4 count.

# Kidney diseases associated with HIV

**HIVAN** 

HIVICK

#### Non-HIVAN

- Non-collapsing FSGS
- TMA
- Amyloidosis (AA) skin poppers
- Crystalline tubulopathy
- Immunotactoid glomerulopathy

#### **HIVAN**

#### First Described

1984 from New York University

#### **Imaging Findings**

Enlarged kidneys

#### **Collapsing FSGS**

- Microcystic dilatation
- Interstitial inflammation
- Endothelial tubuloreticular inclusions

#### **HIV Location**

Could be found in the podocytes and renal tubular epithelial cells

Treatment with cART is effective at reversing HIVAN

#### Clinically

- Heavy proteinuria
- Rapid progression to ESKD
- Almost exclusively in African (Americans)

#### **HIVICK**

#### Lupus-like GN

- Mesangial and endocapillary hypercellularity
- Full-house patten

#### Other GN

 MN, IgAN, PIGN, MPGN, MesPGN, Cryo GN.

#### **EM Findings**

- Immune complex by EM
- Tubuloreticular bodies

#### **Laboratory Findings**

- Negative lupus serologies
- Hypergammaglobulinemia
- Hypocomplementermia
- Hematuria
- Proteinuria (less than HIVAN).

#### **KDIGO** recommend

Term HIVICK gets replaced with a specific description of the pattern of immune complex disease "in the setting of HIV.

#### Risk factors associated with HIVAN and HIVICK

#### **HIVAN**

- Younger
- African
  - APOL1 (G1 and G2) risk alleles (75% have both alleles)
- CART naïve
  - Low CD4 (<200/ml)</li>
  - High viral load
- Develops early in the HIV infection

#### HIVICK

- Older
- Non-Africans outside of Africa
- CART (5-fold higher)
  - Lower viral load
  - Higher CD4 count
- Develops late in HIV infection
- More common than HIVAN since cART

#### **Treatment**

#### **HIVAN**

- CART
- RAS inhibition
  - Prevents ESKD
  - Reverses AKI
  - May prevent the development of HIVAN
- Glucocorticoids\*
  - IDSA recommends a trial of steroids
  - UpToDate does not recommend routine use
  - Lower creatinine and proteinuria
  - Increase infection
  - Relapse after discontinuation

#### **HIVICK**

- Benefits unclear
  - CART
  - RAS blockade
  - Immunosuppression
  - Treat other infections (hepatitis B, C)

## HBV associated kidney diseases

#### **Prevalence**

3-20% of chronic hepatitis B patients may develop HBV-GN

#### Types of Glomerulonephritis

- Membranous nephropathy
- MPGN
- Mes PGN
- IgA nephropathy
- Cryoglobulinemia
- Polyarteritis nodosa

# **Pathogenesis**

#### **Endemic Areas**

More common in endemic areas

Male children

#### **Genotype Role**

 Genotype A is more common in MN and MPGN

# **Chronic Hepatitis Infection**

- Hepatitis B antigen-antibody complexes
- Hepatitis B surface and core antigen can be found on GBM and mesangium.

#### **Membranous Nephropathy**

- Seroconversion from HBeAg to anti-HBe
- Spontaneous remission is uncommon in adults
- Anti-PLA2R and THSD7A are positive in HBV-MN patients probably in the same proportion as non-HBV infected patients

## **KDIGO, 2021**

- Patients with HBV (DNA levels >2000 IU/ml) & GN receive nucleos(t)ide analogues as recommended as per HBV treatment.(1C)
- Pegylated interferon should not be used to treat patients with replicative HBV infection and GN
- **Rituximab and cyclophosphamide** should be **avoided** in patients with simultaneous HBV infection and anti-PLA2R antibody-mediated MN **until a sustained virologic remission** has been obtained by nucleost)ide analogue therapy as it may accelerate HBV replication.

#### Note:

- Plasma exchange may be tried in patients with accompanying cryoglobulinemic vasculitis
- HBV Vaccination
  - Ideally all individuals should be vaccinated

# **HCV-Associated Kidney Diseases**

#### Cryoglobulinemic GN

- Diffuse mesangial proliferative
- MPGN

Membranous nephropathy

#### **Others**

- Fibrillary glomerulonephritis
- Focal segmental glomerulosclerosis
- IgA nephropathy
- Immunotactoid glomerulopathy
- PAN\*

#### **Treatment of HCV-Associated GN**

#### DAA

 Complete remission has been achieved by sofosbuvir and daclatasvir

#### Immunosuppressive therapy

To start 1 - 4 months prior to antiviral therapy

- Plasma exchange
- Cyclophosphamide
- Mycophenolate mofetil
- Glucocorticoid
- RAS blockade\*

#### Rituximab


#### For cryoglobulinemic vasculitis

- Superior to immunosuppressive therapy
- Effective in patients who failed antiviral therapy
- Doubles the effectiveness of antiviral therapy alone

#### Schistosomiasis associated GN

Immune response can start once the egg deposited after infection. Pathology: mesangial proliferative glomerulonephritis. Clinically: proteinuria, hematuria, & sometimes nephrotic syndrome. **Treatment:**  Antiparasitic medications for underlying schistosomiasis Supportive care for associated kidney disease

#### Malaria associated GN





## Conclusion

- Despite global advancements in healthcare, IRGN remains prevalent in regions with high rates of infection and limited access to healthcare.
- The evolving spectrum of IRGN has shifted towards non-streptococcal and adult-onset cases, requiring heightened vigilance and tailored management.
- Early recognition and appropriate treatment of underlying infections are critical to improving patient outcomes.
- Prevention strategies, such as vaccination and improved sanitation, can significantly reduce the disease burden.
- Ongoing research into pathophysiology and biomarkers will pave the way for targeted therapies.

EASED TO INVITE YOU TO OUR NEXT ANNUAL ANNUAL AND

NOVEMBER 27-29th, 2025. Sohar, Sultanate of oman

# 3rd OSNT Annual International Nephrology Conference

**ENDORSED BY** 









